arXiv:1402.0668v1 [math.CO] 4 Feb 2014

An Erdős-Ko-Rado theorem for permutations with fixed number of cycles

Cheng Yeaw Ku * Kok Bin Wong †

February 5, 2014

Abstract

Let S_n denote the set of permutations of $[n] = \{1, 2, \ldots, n\}$. For a positive integer k, define $S_{n,k}$ to be the set of all permutations of $[n]$ with exactly k disjoint cycles, i.e.,

$$S_{n,k} = \{\pi \in S_n : \pi = c_1 c_2 \cdots c_k\},$$

where $c_1, c_2, \ldots, c_k$ are disjoint cycles. The size of $S_{n,k}$ is given by $\begin{bmatrix} n \\ k \end{bmatrix} = (-1)^{n-k} s(n,k)$, where $s(n,k)$ is the Stirling number of the first kind. A family $\mathcal{A} \subseteq S_{n,k}$ is said to be *t-intersecting* if any two elements of $\mathcal{A}$ have at least t common cycles. In this paper, we show that, given any positive integers k, t with $k \geq t+1$, there exists an integer $n_0 = n_0(k,t)$, such that for all $n \geq n_0$, if $\mathcal{A} \subseteq S_{n,k}$ is t-intersecting, then

$$|\mathcal{A}| \leq \begin{bmatrix} n-t \\ k-t \end{bmatrix},$$

with equality if and only if $\mathcal{A}$ is the stabiliser of t fixed points.

KEYWORDS: t-intersecting family, Erdős-Ko-Rado, permutations

1 Introduction

Let $[n] = \{1, \ldots, n\}$, and let $\binom{[n]}{k}$ denote the family of all k-subsets of $[n]$. A family $\mathcal{A}$ of subsets of $[n]$ is *t-intersecting* if $|A \cap B| \geq t$ for all $A, B \in \mathcal{A}$. One of the most beautiful results in extremal combinatorics is the Erdős-Ko-Rado theorem.

Theorem 1.1 (Erdős, Ko, and Rado [13], Frankl [14], Wilson [37]). *Suppose $\mathcal{A} \subseteq \binom{[n]}{k}$ is t-intersecting and $n > 2k - t$. Then for $n \geq (k-t+1)(t+1)$, we have*

$$|\mathcal{A}| \leq \binom{n-t}{k-t}.$$

Moreover, if $n > (k-t+1)(t+1)$ then equality holds if and only if $\mathcal{A} = \{A \in \binom{[n]}{k} : T \subseteq A\}$ for some t-set T.

*Department of Mathematics, National University of Singapore, Singapore 117543. E-mail: matkcy@nus.edu.sg

†Institute of Mathematical Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia. E-mail: kbwong@um.edu.my.

Later, Ahlswede and Khachatrian [1] extended the Erdős-Ko-Rado theorem by determining the structure of all t-intersecting set systems of maximum size for all possible n (see also [3, 15, 22, 28, 32, 34, 35] for some related results). There have been many recent results showing that a version of the Erdős-Ko-Rado theorem holds for combinatorial objects other than set systems (see [2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 29, 30, 31, 33, 36, 38]). Most notably is the results of Ellis, Friedgut and Pilpel [12] who showed that for sufficiently large n depending on t, a t-intersecting family $\mathcal{A}$ of permutations has size at most $(n-t)!$, with equality if and only if $\mathcal{A}$ is a coset of the stabilizer of t points, thus settling an old conjecture of Deza and Frankl in the affirmative. The proof uses spectral methods and representations of the symmetric group.

Let S_n denote the set of permutations of $[n]$. For a positive integer k, define $S_{n,k}$ to be the set of all permutations of $[n]$ with exactly k disjoint cycles, i.e.,

$$S_{n,k} = \{\pi \in S_n : \pi = c_1 c_2 \cdots c_k\},$$

where $c_1, c_2, \ldots, c_k$ are disjoint cycles. It is well known that the size of $S_{n,k}$ is given by $\begin{bmatrix} n \\ k \end{bmatrix} = (-1)^{n-k} s(n,k)$, where $s(n,k)$ is the *Stirling number of the first kind*.

We shall use the following notations:

(a) $N(c) = \{a_1, a_2, \ldots, a_l\}$ for a cycle $c = (a_1, a_2, \ldots, a_l)$;

(b) $M(\pi) = \{c_1, c_2, \ldots, c_k\}$ for a $\pi = c_1 c_2 \ldots c_k \in S_{n,k}$;

A family $\mathcal{A} \subseteq S_{n,k}$ is said to be *t-intersecting* if any two elements of $\mathcal{A}$ have at least t common cycles, i.e., $|M(\pi_1) \cap M(\pi_2)| \geq t$ for all $\pi_1, \pi_2 \in \mathcal{A}$.

Theorem 1.2. *Given any positive integers k, t with $k \geq t+1$, there exists an integer $n_0 = n_0(k,t)$, such that for all $n \geq n_0$, if $\mathcal{A} \subseteq S_{n,k}$ is t-intersecting, then*

$$|\mathcal{A}| \leq \begin{bmatrix} n-t \\ k-t \end{bmatrix},$$

with equality if and only if $\mathcal{A}$ is the stabiliser of t fixed points.

2 Stirling number revisited

The unsigned Stirling number $\begin{bmatrix} n \\ k \end{bmatrix}$ satisfies the recurrence relation

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix} + (n-1) \begin{bmatrix} n-1 \\ k \end{bmatrix}, \tag{1}$$

with initial conditions $\begin{bmatrix} 0 \\ 0 \end{bmatrix} = 1$ and $\begin{bmatrix} n \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ k \end{bmatrix} = 0$, $n > 0$. Note that $\begin{bmatrix} n \\ n \end{bmatrix} = 1$. By using equation (1) and induction on n,

$$\begin{bmatrix} n \\ 1 \end{bmatrix} = (n-1)!. \tag{2}$$

By applying equation (1) repeatedly,

$$\begin{aligned}
\begin{bmatrix} n \\ k \end{bmatrix} &= \begin{bmatrix} n-1 \\ k-1 \end{bmatrix} + (n-1)\begin{bmatrix} n-1 \\ k \end{bmatrix} \\
&= \begin{bmatrix} n-1 \\ k-1 \end{bmatrix} + (n-1)\begin{bmatrix} n-2 \\ k-1 \end{bmatrix} + (n-1)(n-2)\begin{bmatrix} n-2 \\ k \end{bmatrix} \\
&= \begin{bmatrix} n-1 \\ k-1 \end{bmatrix} + (n-1)\begin{bmatrix} n-2 \\ k-1 \end{bmatrix} + (n-1)(n-2)\begin{bmatrix} n-3 \\ k-1 \end{bmatrix} + (n-1)(n-2)(n-3)\begin{bmatrix} n-3 \\ k \end{bmatrix} \\
&\vdots \\
&= \sum_{r=k-1}^{n-1} \frac{(n-1)!}{r!}\begin{bmatrix} r \\ k-1 \end{bmatrix}. \qquad (3)
\end{aligned}$$

In particular (by equations (2) and (3)),

$$\begin{bmatrix} n \\ 2 \end{bmatrix} = (n-1)!\sum_{r=1}^{n-1}\frac{1}{r}. \qquad (4)$$

By elementary calculus, it is easy to show that for sufficiently large n,

$$\frac{\ln n}{2} < \ln(n-1) + \frac{1}{n-1} \le \sum_{r=1}^{n-1}\frac{1}{r} \le \ln(n-1) + 1 < 2\ln n. \qquad (5)$$

Hence, there are positive constants $\alpha(2)$ and $\beta(2)$ such that

$$\alpha(2)((n-1)!(\ln n)) < \begin{bmatrix} n \\ 2 \end{bmatrix} < \beta(2)((n-1)!(\ln n)), \qquad (6)$$

for all $n \ge 2$.

Again, by elementary calculus, for $m \ge 1$ and sufficiently large n depending on m,

$$\frac{\ln^{m+1} n}{2(m+1)} < \sum_{r=1}^{n-1}\frac{\ln^m r}{r} < \frac{2\ln^{m+1} n}{m+1}. \qquad (7)$$

The following lemma follows by using equations (3) and (7), and induction on k.

Lemma 2.1. *There are positive constants $\alpha(k)$ and $\beta(k)$ such that*

$$\alpha(k)((n-1)!(\ln^{k-1} n)) < \begin{bmatrix} n \\ k \end{bmatrix} < \beta(k)((n-1)!(\ln^{k-1} n)),$$

for all $n \ge k$.

3 Main results

A family $\mathcal{B} \subseteq S_{n,k}$ is said to be *independent* if $M(\pi_1) \cap M(\pi_2) = \varnothing$ for all $\pi_1, \pi_2 \in \mathcal{B}$.

Lemma 3.1. *Let $\mathcal{A} \subseteq S_{n,k}$ and $k \geq 2$. If a maximal independent subset of $\mathcal{A}$ is of size at most l, then*

$$|\mathcal{A}| \leq kl \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}.$$

Proof. Let $\mathcal{B} = \{\pi_1, \pi_2, \ldots, \pi_l\}$ be a maximal independent subset of $\mathcal{A}$ of size l. Let $\pi_i = c_{i1}c_{i2}\ldots c_{ik}$ where $c_{i1}, c_{i2}, \ldots, c_{ik}$ are disjoint cycles, and

$$Q = \bigcup_{i=1}^{l} M(\pi_i).$$

Note that $|Q| = kl$.

Let

$$\mathcal{A}_{ij} = \{\pi \in \mathcal{A} \ : \ c_{ij} \in M(\pi)\}.$$

Let $\pi \in \mathcal{A} \setminus \mathcal{B}$. By the maximality of $\mathcal{B}$, $M(\pi) \cap Q \neq \varnothing$. So, $c_{ij} \in M(\pi)$ for some i, j, and $\pi \in \mathcal{A}_{ij}$. Hence,

$$\mathcal{A} = \bigcup_{i,j} \mathcal{A}_{ij}.$$

The lemma follows by noting that

$$|A_{ij}| \leq \begin{bmatrix} n - |N(c_{ij})| \\ k-1 \end{bmatrix} \leq \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}.$$

□

Let $\mathcal{A} \subseteq S_{n,k}$ and $c_1, \ldots, c_t$ by cycles such that $N(c_i) \cap N(c_j) = \varnothing$ for $i \neq j$. Let $T = \{c_1, \ldots, c_t\}$. We set

$$\mathcal{A}(T) = \{\pi \in \mathcal{A} \ : \ T \subseteq M(\pi)\}.$$

Now, for each element $\pi \in \mathcal{A}(T)$, we remove all the cycles $c_1, c_2, \ldots, c_t$ from π and denote the resulting set by $\mathcal{A}^*(T)$. Let $P = \bigcup_{i=1}^{t} N(c_i)$. Note that $|\mathcal{A}(T)| = |\mathcal{A}^*(T)|$ and $\mathcal{A}^*(T) \subseteq S_{n-|P|,k-t}$. Here, $S_{n-|P|,k-t}$ is the set of all permutations of $[n] \setminus P$ with exactly $k-t$ disjoint cycles.

Lemma 3.2. *Let $\mathcal{A} \subseteq S_{n,k}$ be maximal t-intersecting and $k \geq t+1$. Let $T = \{c_1, \ldots, c_t\}$ with $N(c_i) \cap N(c_j) = \varnothing$ for $i \neq j$. If $\mathcal{A}^*(T)$ has an independent set of size at least $k+1$, then*

$$\mathcal{A} = \{\pi \in S_{n,k} \ : \ T \subseteq M(\pi)\}.$$

Proof. Let $\{\pi_1, \ldots, \pi_{k+1}\}$ be an independent subset of $\mathcal{A}^*(T)$ of size $k+1$. For $l = 1, 2, \ldots, k+1$, let

$$\pi_l = c_1 \ldots c_t d_{l,t+1} \ldots d_{l,k},$$

where $c_1, \ldots, c_t, d_{l,t+1}, \ldots, d_{l,k}$ are disjoint cycles.

Suppose there is a $\pi \in \mathcal{A}$ such that $c_{i_0} \notin M(\pi)$ for a fixed i_0. Since $\mathcal{A}$ is t-intersecting,

$$|M(\pi) \cap M(\pi_l)| \geq t.$$

Therefore there is a j_l $(t+1 \leq j_l \leq k)$ with $d_{l,j_l} \in M(\pi)$. Since all the d_{l,j_l} are distinct, $k = |M(\pi)| \geq k+1$, a contradiction. Hence,

$$\mathcal{A} \subseteq \{\pi \in S_{n,k} \ : \ T \subseteq M(\pi)\}.$$

By the maximality of $\mathcal{A}$, the lemma follows. □

Proof of Theorem 1.2. We may assume that $\mathcal{A}$ is maximal t-intersecting.

Suppose $k = t+1$. Since $\mathcal{A}$ is t-intersecting, there are $\pi_1, \pi_2 \in \mathcal{A}$ such that

$$\pi_1 = c_1 c_2 \dots c_t d_1$$
$$\pi_2 = c_1 c_2 \dots c_t d_2$$

where $c_1, \dots, c_t, d_1$ are disjoint cycles, $d_2 \neq d_1$ and $N(d_2) = N(d_1)$. Suppose there is a $\pi \in \mathcal{A}$ with $c_{i_0} \notin M(\pi)$ for some i_0. Then $d_1, d_2 \in M(\pi)$. But this is impossible as $N(d_1) = N(d_2)$. Hence,

$$\mathcal{A} = \{\pi \in S_{n,k} \ : \ c_i \in M(\pi) \ \text{ for } \ i = 1, 2, \dots, t\},$$

for $\mathcal{A}$ is maximal t-intersecting. Let $P = \bigcup_{i=1}^{t} N(c_i)$. Then

$$|\mathcal{A}| = \begin{bmatrix} n - |P| \\ 1 \end{bmatrix} \leq \begin{bmatrix} n-t \\ 1 \end{bmatrix},$$

with equality if and only if $|N(c_i)| = 1$ for $i = 1, 2 \dots, t$, i.e., $\mathcal{A}$ is the stabilizer of at least t fixed points. Now, if $n \geq t+2$, then $\mathcal{A}$ is the stabilizer of t fixed points.

Suppose $k \geq t+2$. Let $\pi_0 = d_1 d_2 \dots d_k \in \mathcal{A}$ be fixed, where $d_1, \dots, d_k$ are disjoint cycles. Then

$$\mathcal{A} = \bigcup_{T \subseteq M(\pi_0), |T| = t} \mathcal{A}(T).$$

Case 1. Suppose that for each $T \subseteq M(\pi_0)$ with $|T| = t$, all independent subsets of $\mathcal{A}^*(T)$ is of size at most k. Then by Lemma 3.1 and equation (1),

$$|\mathcal{A}(T)| = |\mathcal{A}^*| \leq k^2 \begin{bmatrix} n - |P| - 1 \\ k - t - 1 \end{bmatrix} \leq k^2 \begin{bmatrix} n-t-1 \\ k-t-1 \end{bmatrix},$$

where $P = \bigcup_{c \in T} N(c)$. This implies that

$$|\mathcal{A}| \leq k^2 \binom{k}{t} \begin{bmatrix} n-t-1 \\ k-t-1 \end{bmatrix}.$$

By Lemma 2.1, there is are positive constants α and β such that

$$|\mathcal{A}| < \beta k^2 \binom{k}{t} \left((n-t-2)! (\ln^{k-t-2} n) \right),$$

and

$$\alpha \left((n-t-2)! (\ln^{k-t-1} n) \right) < \begin{bmatrix} n-t-1 \\ k-t \end{bmatrix}.$$

So, for sufficiently large n, $|\mathcal{A}| < \begin{bmatrix} n-t-1 \\ k-t \end{bmatrix}$, and by equation (1), $|\mathcal{A}| < \begin{bmatrix} n-t \\ k-t \end{bmatrix}$.

Case 2. Suppose that there is a $T_0 \subseteq M(\pi_0)$ with $|T_0| = t$, such that $\mathcal{A}^*(T_0)$ has an independent set of size at least $k+1$. By Lemma 3.2,

$$\mathcal{A} = \{\pi \in S_{n,k} \ : \ T_0 \subseteq M(\pi)\}.$$

Let $P_0 = \bigcup_{c \in T_0} N(c_0)$. Then

$$|\mathcal{A}| = \begin{bmatrix} n-|P_0| \\ k-t \end{bmatrix} \leq \begin{bmatrix} n-t \\ k-t \end{bmatrix},$$

with equality if and only if $|N(c_i)| = 1$ for $i = 1, 2 \ldots, t$, i.e., $\mathcal{A}$ is the stabilizer of at least t fixed points. □

Acknowledgments

This project is supported by the Advanced Fundamental Research Cluster, University of Malaya (UMRG RG238/12AFR).

References

[1] R. Ahlswede and L. H. Khachatrian, The complete intersection theorem for systems of finite sets, *European J. Combin.* **18** (1997), 125–136.

[2] R. Ahlswede and L.H. Khachatrian, The diametric theorem in Hamming spaces – Optimal anticodes, *Adv. in Appl. Math.* **20** (1998), 429–449.

[3] C. Bey, On cross-intersecting families of sets, *Graphs Combin.* **21** (2005), 161–168.

[4] P. Borg, Extremal t-intersecting sub-families of hereditary families, *J. London Math. Soc.* **79** (2009), 167–185.

[5] P. Borg, On t-intersecting families of signed sets and permutations, Discrete Math. **309** (2009), 3310–3317.

[6] P. Borg and F.C. Holroyd, The Erdos-Ko-Rado property of various graphs containing singletons, *Discrete Math.* **309** (2009), 2877–2885.

[7] F. Brunk and S. Huczynska, Some Erdos-Ko-Rado theorems for injections, *European J. Combin.* **31** (2010), 839–860.

[8] P. J. Cameron and C. Y. Ku, Intersecting families of permutations, *European J. Combin.* **24** (2003), 881–890.

[9] A. Chowdhury and B. Patkós, Shadows and intersections in vector spaces, *J. Combin. Theory Ser. A* **117** (2010), 1095–1106.

[10] M. Deza and P. Frankl, On the maximum number of permutations with given maximal or minimal distance, *J. Combin. Theory Ser. A* **22** (1977), 352–360.

[11] D. Ellis, Stability for t-intersecting families of permutations, *J. Combin. Theory Ser. A* **118** (2011), 208–227.

[12] D. Ellis, E. Friedgut and H. Pilpel, Intersecting families of permutations, *Journal of the American Society* **24** (2011), 649-682.

[13] P. Erdős, C. Ko and R. Rado, Intersection theorems for systems of finite sets, *Quart. J. Math. Oxford Ser.* 2 **12** (1961), 313–318.

[14] P. Frankl, The Erdős-Ko-Rado theorem is true for $n = ckt$, *Col. Soc. Math. J. Bolyai* **18** (1978), 365–375.

[15] P. Frankl and N. Tokushige, On r-cross intersecting families of sets, *Combin. Probab. Comput.* **20** (2011), 749–752.

[16] P. Frankl and R.M. Wilson, The Erdős-Ko-Rado theorem for vector spaces, *J. Combin. Theory Ser. A* **43** (1986) 228–236.

[17] C. Godsil and K. Meagher, A new proof of the Erdős-Ko-Rado theorem for intersecting families of permutations, *European J. Combin.* **30** (2009), 404–414.

[18] A.J.W. Hilton and C.L. Spencer, A graph-theoretical generalisation of Berges analogue of the Erdos-Ko-Rado theorem, Trends in Graph Theory, Birkhauser Verlag, Basel, Switzerland (2006), 225–242.

[19] F.C. Holroyd, C. Spencer and J. Talbot, Compression and Erdos-Ko-Rado graphs, *Discrete Math.* **293** (2005), 155–164.

[20] F.C. Holroyd and J. Talbot, Graphs with the Erdos-Ko-Rado property, *Discrete Math.* **293** (2005), 165–176.

[21] G. Hurlbert and V. Kamat, Erdős-Ko-Rado theorems for chordal graphs and trees, *J. Combin. Theory Ser. A* **118** (2011), 829–841.

[22] P. Keevash, Shadows and intersections: Stability and new proofs, Adv. Math. **218** (2008) 1685–1703.

[23] C. Y. Ku and I. Leader, An Erdős-Ko-Rado theorem for partial permutations, *Discrete Math.* **306** (2006), 74–86.

[24] C. Y. Ku and D. Renshaw, Erdős-Ko-Rado theorems for permutations and set partitions, *J. Combin. Theory Ser. A* **115** (2008), 1008–1020.

[25] C. Y. Ku and T. W. H. Wong, Intersecting families in the alternating group and direct product of symmetric groups, *Electron. J. Combin.* **14** (2007), #R25.

[26] C. Y. Ku and K. B. Wong, An analogue of Hilton-Milner theorem for set partitions, *J. Combin. Theory Ser. A* **120** (2013), 1508–1520.

[27] C. Y. Ku and K. B. Wong, On cross-intersecting families of set partitions, *Electron. J. Combin.* **19**(4) (2012), #49.

[28] C. Y. Ku and K. B. Wong, On r-cross intersecting families of sets, *Far East J. Math. Sci.* **75** (2013), 295–300.

[29] C. Y. Ku and K. B. Wong, An analogue of the Erdős-Ko-Rado theorem for weak compositions, *Discrete Math.* **313** (2013) 2463-2468.

[30] B. Larose and C. Malvenuto, Stable sets of maximal size in Kneser-type graphs, *European J. Combin.* **25** (2004), 657–673.

[31] Y.-S. Li and Jun Wang, Erdos-Ko-Rado-type theorems for colored sets, *Electron. J. Combin.* **14** (2007) #R1.

[32] M. Matsumoto and N. Tokushige, The exact bound in the Erdős-Ko-Rado theorem for cross-intersecting families, *J. Combin. Theory Ser. A* **52** (1989), 90–97.

[33] A. Moon, An analogue of the Erdős-Ko-Rado theorem for the Hamming schemes $H(n, q)$, *J. Combin. Theory Ser. A* **32**(3) (1982), 386–390.

[34] L. Pyber, A new generalization of the Erdős-Ko-Rado theorem, *J. Combin. Theory Ser. A* **43** (1986), 85–90.

[35] N. Tokushige, A product version of the Erdős-Ko-Rado theorem, *J. Combin. Theory Ser. A* **118** (2011), 1575–1587.

[36] J. Wang and S. J. Zhang, An Erdős-Ko-Rado-type theorem in Coxeter groups, *European J. Combin.* **29** (2008), 1112–1115.

[37] R.M. Wilson, The exact bound in the Erdős-Ko-Rado theorem, *Combinatorica* **4** (1984), 247–257.

[38] R. Woodroofe, Erdos-Ko-Rado theorems for simplicial complexes, *J. Combin. Theory Ser. A* **118** (2011), 1218–1227.